

Growth Hormone Induces Low-Density Lipoprotein Clearance but not Bile Acid Synthesis in Humans

Suzanne Lind, Mats Rudling, Sverker Ericsson, Hans Olivecrona, Mats Eriksson, Birgit Borgström, Gösta Eggertsen, Lars Berglund, Bo Angelin

Objective—Growth hormone (GH) induces hepatic low-density lipoprotein (LDL) receptors and lowers plasma cholesterol. We characterized the influence of GH treatment on plasma LDL clearance in normal humans and investigated the relative role of LDL receptor (LDLR) activity and stimulation of bile acid synthesis in subjects with different LDLR expression.

Methods and Results—Plasma clearance of autologous ^{125}I -LDL was measured before and during 3 weeks of treatment with GH (0.1 IU/kg per day) in 9 healthy young males. Plasma LDL cholesterol was reduced by 13% and the fractional catabolic rate of LDL increased by 27%. More marked changes were seen in a patient with hypopituitarism substituted with GH (0.07 IU/kg per day) for 3 months. In a second study, GH dose-dependently reduced LDL cholesterol and increased Lp(a) levels in 3 groups of males: younger and elderly healthy subjects and heterozygous familial hypercholesterolemia (FH). No effect on bile acid synthesis measured by the plasma marker 7α -hydroxy-4-cholesten-3-one was observed. In an LDLR-deficient FH homozygote, LDL cholesterol was not affected by GH.

Conclusions—GH treatment reduces plasma LDL cholesterol by inducing LDL clearance. In humans, LDLR expression is a prerequisite for this effect, whereas it is not related to stimulation of bile acid synthesis. (*Arterioscler Thromb Vasc Biol.* 2004;24:349-356.)

Key Words: insulin-like growth factor-I ■ low density lipoprotein kinetics
■ familial hypercholesterolemia ■ lipoprotein (a)

Increased levels of plasma low-density lipoprotein (LDL) cholesterol are associated with an enhanced risk of atherosclerosis and coronary heart disease. LDL receptors (LDLRs) are of great importance in determining plasma cholesterol, as evident from the genetic disease, familial hypercholesterolemia (FH), in which LDLRs are reduced by $\approx 50\%$ (heterozygotes) or lost (homozygotes).¹ Stimulation of LDLR activity in the liver is a major mechanism for reducing plasma LDL cholesterol levels in humans^{2,3} and may be achieved by inhibition of cholesterol biosynthesis or stimulation of bile acid production.^{4–6} Also, estrogen treatment markedly induces hepatic LDLRs.⁷ In all these situations, the plasma clearance of LDL is enhanced, whereas the LDL production may show a more variable response.^{8–11}

We have previously reported that treatment of normal adults with growth hormone (GH) stimulates hepatic LDLRs to a magnitude similar to that seen during therapy with cholesterol synthesis inhibitors.¹² GH treatment can reduce total and LDL cholesterol but increases lipoprotein(a) (Lp(a)) in normal^{13–15} and GH-deficient adults.^{16–18} In parallel to the

reduced secretion of GH with aging in humans,^{14,19} plasma LDL clearance and bile acid synthesis are reduced,^{20,21} whereas plasma LDL cholesterol increases.^{19,22} In the rat, the age-dependent reduction of bile acid production and increase of plasma cholesterol can be reversed by GH treatment.²³ Thus, it has been hypothesized that GH is important for the normal regulation of plasma cholesterol.^{19,24} This may occur through effects of GH on hepatic LDLRs and/or bile acid synthesis.^{23,25} Interestingly, the administration of GH can stimulate bile acid synthesis and reduce plasma cholesterol even in mice lacking LDLRs.²⁶

A more detailed knowledge of the effects of GH on plasma LDL metabolism is therefore warranted. In the present study, we first show that GH stimulates the plasma clearance of LDL apoB, resulting in reduced plasma LDL cholesterol levels, whereas it has minor effects on LDL apoB synthesis. Further, we address the following questions. Is the basal expression of LDLRs important for the LDL-lowering effect of GH in humans? Is the effect of GH in humans related to stimulation of bile acid synthesis? Is the effect of GH on

Received October 3, 2003; revision accepted November 21, 2003.

From the Metabolism Unit (S.L., M.R., S.E., H.O., M.E., B.A.), Center for Metabolism and Endocrinology, Department of Medicine and Molecular Nutrition Unit, Center for Nutrition and Toxicology, Novum; Division of Clinical Chemistry (S.L., G.E., L.B.), Department of Laboratory Medicine; and Pediatric Unit (B.B.), Department of Clinical Sciences, Karolinska Institute at Huddinge University Hospital, Stockholm, Sweden.

S.L. and S.E. contributed equally to this work.

Correspondence to Bo Angelin, Center for Metabolism & Endocrinology, M63 Huddinge University Hospital, S-141 86 Stockholm, Sweden. E-mail bo.angelin@medhs.ki.se

© 2004 American Heart Association, Inc.

Arterioscler Thromb Vasc Biol. is available at <http://www.atvbaha.org>

DOI: 10.1161/01.ATV.0000110657.67317.90

plasma Lp(a) related to the LDL-lowering effect? We investigated four groups of human subjects in whom the expression of LDLRs can be assumed to differ: (1) normal younger males; (2) normal elderly males; (3) middle-aged males heterozygous for FH; and (4) a homozygote FH child.

Materials and Methods

Subjects

In the LDL turnover study, 9 healthy male volunteers aged 24 to 41 years (mean \pm SEM: 32 \pm 2 years) participated. They were of normal weight, with BMIs ranging from 22.6 to 25.8 (24.2 \pm 0.43). Plasma cholesterol and triglycerides were below the 95th percentile of the Swedish population. There was no clinical or laboratory evidence of thyroid, kidney, heart, or liver disease, diabetes, or alcohol abuse; no medication was used. In addition, 1 male patient (age: 61 years, BMI: 30.6) with hypopituitarism after surgical removal of a non-secreting adenoma 2 years before the study was investigated. He received substitution therapy with testosterone, thyroid hormone, and cortisone.

For the dose-response study, 8 men with heterozygous FH (age: 45 \pm 5 years; BMI: 25.0 \pm 0.95) and 15 healthy male volunteers were studied. The diagnosis of FH was based on described criteria.^{27,28} Five patients had known mutations in the LDLR gene,^{28,29} two had FH-Helsinki, one had a deletion of exon 2-3, and 2 had point mutations (L393R, IVS 10-2 A>G). Lipid-lowering drugs were omitted 6 weeks before the study. The healthy males were recruited from 2 age groups: younger (28 \pm 1; range 23 to 33 years) and elderly (66 \pm 1; 62 to 70 years) subjects. Neither had clinical or laboratory evidence of diabetes, overweight, hypertension, or renal, thyroid, or hepatic disease; BMIs were 24.4 \pm 1.1 and 26.3 \pm 0.83, respectively.

In addition, one homozygous FH patient of Turkish origin was investigated. She was 6 years old and had characteristic xanthomas and plasma cholesterol of \approx 20 mmol/L without treatment. DNA analysis confirmed homozygosity for the LDLR gene mutation W556R, which has been reported to result in <5% of normal LDLR activity in cultured fibroblasts.³⁰ Her apoE genotype was apoE3/2. Although she had increased intima media thickness of the carotids, echocardiography and workload test results were normal.

Informed consent was obtained from all participants and from the parents of the child. The studies were approved by the Ethics Committee of the Karolinska Institute and by the Swedish Medical Products Agency.

Experimental Procedures

In the LDL turnover study, the subjects were studied twice: once in the basal state and once during GH treatment. They were followed-up as outpatients and given a standardized diet 5 days before and during each study period.^{10,20} Potassium iodide, 200 mg daily, was administered orally 5 days before and during the studies. Recombinant human GH (Genotropin; Pharmacia, Sweden), 0.1 IU/kg per day, was administered for 3 weeks as single subcutaneous injections at 8:00AM. Plasma LDLs were isolated, labeled with ¹²⁵I, and re-injected into the subject within 5 days of sampling; during the second period, plasma was obtained immediately before administration of the first dose of GH. Fasting blood samples were analyzed repeatedly for glucose, plasma cholesterol and triglycerides, lipoprotein lipids, insulin, insulin-like growth factor-I (IGF-I), and IGF binding protein-1. Fasting blood samples were obtained just before the daily GH injection. Constant body weight and plasma lipoprotein levels indicated a metabolic steady state during the studies. The patient with hypopituitarism was studied using a similar protocol (except during the second period, when LDLs were isolated during ongoing treatment) before and after 3 months of therapy, with GH administered at a daily dose of 0.07 IU/kg per day.

In the dose-response study, the 3 groups (younger, elderly, and FH heterozygotes) were administered 3 doses (0.025, 0.05, and 0.1 IU/kg per day) of GH for 1 week each for a total time of 3 weeks. GH was injected subcutaneously at 8:00PM daily. Blood sampling was performed every 7 days in the morning after overnight fast, when

blood pressure, body weight, biochemical safety tests, and side effects were monitored.

The homozygous FH child had an individual protocol. Initially, she was administered continued treatment with 0.6 mg/kg per day of atorvastatin (Lipitor; Pfizer, Sweden). She was first administered GH 0.1 IU/kg per day for 1 week in addition to atorvastatin; thereafter, both drugs were omitted for a 4-week washout period. GH (0.1 IU/kg per day) was administered for 1 week and then increased to 0.2 IU/kg per day. Colestipol 0.45 g/kg per day (Lestid; Pharmacia, Sweden) and atorvastatin 0.6 mg/kg per day were then added to GH treatment as described in Figure 5.

Assays

Plasma insulin was determined by radioimmunoassay (Pharmacia & Upjohn Diagnostics AB, Uppsala, Sweden) and serum IGF-1 by RIA.³¹ The concentrations of IGF binding protein-1 were determined according to Póvoa et al³² (courtesy of Prof Kerstin Hall, Karolinska Institutet).

Plasma cholesterol and triglycerides were measured by enzymatic techniques (Boehringer-Mannheim, Mannheim, Germany), and lipoproteins were quantitated using a combination of ultracentrifugation and precipitation.^{33,34} For the analysis of Lp(a) and apoAI and apoB, immunoturbidimetric methods were used (DAKO A/S, Glostrup, Denmark). The level of LDL apoprotein (LDL apoB) was calculated by multiplying the LDL cholesterol level with the protein/cholesterol ratio in the isolated LDL.¹⁰ Plasma 7 α -hydroxy-4-cholesten-3-one (C4) was analyzed as described,³⁵ and unesterified lathosterol in plasma was quantitated by mass spectrometry.³⁶

¹²⁵I-LDL Preparation and Turnover

LDLs (density 1.02 to 1.063 g/mL) were prepared by sequential ultracentrifugation and labeled with ¹²⁵I using the iodine monochloride method;³⁷ 30 to 60 μ Ci (1.1 to 2.2 MBq) of ¹²⁵I-LDL apoB was injected exactly as described.¹⁰ Blood samples were collected at 10 minutes and at 2, 4, 6, 8, 10, 12, 24, and 36 hours after the injection. Thereafter, samples were collected daily for 14 days. Urine was collected in 24-hour portions during the whole study. The fractional catabolic rate (FCR) of ¹²⁵I-LDL apoB was calculated from the slope of the plasma radioactivity decay curve using the two-compartment model of Matthews.³⁸ In addition, FCR was also calculated independently from the urine/plasma (U/P) ratio.¹⁰ The absolute catabolic rate of LDL apoB, expressed as milligrams of apoprotein synthesized per day normalized for body weight, was estimated according to Langer et al³⁹ as described.^{10,40}

Statistical Analysis

Data are presented as mean \pm SEM or median (range). All parameters were analyzed for normal distribution before and after log transformation. The significance of differences between groups was tested by two-way repeated-measures ANOVA, followed by planned comparison of group means according to the method of Tukey. The correlations between IGF-I and Lp(a), LDL, and HDL cholesterol were calculated using Spearman's rank-order correlation. Calculations were performed using Statistica 5.5 (StatSoft Scandinavia AB, Uppsala, Sweden).

Results

Turnover Study

When exogenous human recombinant GH was administered to healthy males, serum IGF-I levels increased \approx 3-fold after 1 week and remained elevated during the study period (Figure 1). There were no significant changes in blood glucose, serum insulin, or IGF binding protein-1 levels. The levels of IGF-I reached were clearly supranormal and comparable to those observed in patients with mild acromegaly.⁴¹

After 1 week of therapy, a clear and sustained reduction of plasma LDL cholesterol was observed, whereas there were no significant changes in total or HDL cholesterol or in plasma




Figure 1. Changes in blood glucose and serum levels of indicated hormones in response to GH treatment in healthy male volunteers. *** $P < 0.001$ compared with baseline (A and B). Changes in plasma levels of LDL and HDL cholesterol and very-low-density lipoprotein triglycerides in response to GH treatment. * $P < 0.01$ compared with baseline (C).

triglycerides (Figure 1). Also, body weight remained stable during the study, further indicating a steady state. The FCR of LDL apoB averaged 0.315 ± 0.015 pools per day in the basal state as calculated from analyses of plasma radioactivity decay curves (Figure 2). The calculated production rate of LDL apoB was 12.3 ± 1.2 mg/kg per day. Both these values are in good agreement with data obtained previously in healthy males of similar ages.²⁰ Data from urine/protein analyses were in excellent agreement with those obtained from analyses of plasma data (not shown).

A clear increase in the elimination rate of LDL apoB was observed in response to GH treatment (0.399 ± 0.010 pools per day; $P = 0.0023$; Figure 2). Plasma LDL apoB levels were reduced by 14%, from 101 ± 7 to 87 ± 5 mg/dL ($P = 0.037$). The estimated production rate tended to be slightly higher




Figure 2. Individual changes in the kinetics of ¹²⁵I-LDL apo B before (B) and during (D) treatment. The means are indicated by horizontal bars.




Figure 3. Data from the turnover studies with ¹²⁵I-LDL in the hypopituitary patient before and 3 months after (during) the onset of addition of GH to substitution therapy.

during therapy, but the difference was not statistically significant (13.3 ± 1.2 mg/kg per day; $P = 0.14$).

It should be noted that the tracer used in the experiments consisted of LDL collected before GH treatment was initiated. Analyses of the LDL composition (mg%) before (cholesterol:triglyceride:phospholipid:protein = $40 \pm 0.7:7 \pm 0.7:29 \pm 0.7:24 \pm 0.7$) and after 3 weeks of GH treatment ($39 \pm 0.7:10 \pm 1.0:28 \pm 0.7:23 \pm 0.3$) actually revealed an increase in the relative content of triglyceride ($P = 0.002$). The calculations of LDL apoB flux before and during treatment should therefore not be regarded as fully comparable on a quantitative basis. However, the demonstration of an increased clearance of pre-treatment LDL particles in GH-treated subjects is strongly suggestive of an increased hepatic LDLR expression, because the problem of a possible difference in receptor binding affinity of particles isolated during treatment⁴²⁻⁴⁴ is avoided by the present approach.

In a separate experiment (Figure 3), one patient with hypopituitarism and GH deficiency was administered GH at substitution dosage (0.07 IU/kg per day). LDL apoB FCR in this patient increased 38% and the response was even more pronounced than in the normal subjects. Because of an increased production rate, plasma LDL apoB levels were virtually unchanged, whereas mean LDL cholesterol decreased from 4.5 to 3.8 mmol/L.

Dose-Response Study

All participants completed the study with excellent compliance and tolerated the treatment well. In response to GH treatment, plasma levels of IGF-I increased dose-dependently in all groups (Figure 4A). At baseline, younger subjects showed slightly higher IGF-I levels compared with levels of elderly subjects ($P = 0.025$) and of FH patients ($P = 0.05$). They also had a significantly more pronounced response after weeks 2 and 3. Also, plasma insulin and glucose levels increased dose-dependently in all groups (not shown), but no significant differences in responses were seen between groups.

The plasma total and LDL cholesterol levels were reduced by GH treatment in all 3 groups in a dose-dependent way (Table 1). The relative response was somewhat more brisk in




Figure 4. IGF-I levels in all 3 groups during the study (A). The relative changes in plasma LDL cholesterol (B) and Lp(a) (C) levels in the 3 groups expressed as % change of pre-treatment levels. Younger (○); elderly (□); FH patients (●). *P* values significantly different from basal are: **P*<0.05, ***P*<0.01, ****P*<0.001; + indicates significantly different from the other two subject groups (*P*<0.05).

the younger subjects (Figure 4B), but there was no difference between the 3 groups after 3 weeks. HDL cholesterol levels were slightly but significantly reduced in all groups after 1 week, but there was no relationship to dose, and apoAI levels were unaltered. Plasma apoB levels were significantly reduced in all groups after 3 weeks of treatment. FH patients had higher basal triglyceride levels (*P*=0.019) as compared with levels of healthy subjects. Plasma triglycerides increased dose-dependently in all groups. Lp(a) levels increased significantly in all groups in a dose-dependent manner. There were, however, no significant differences in the relative responses between the groups (Figure 4C).

Although the IGF-I response to GH treatment was stronger in the younger subjects (Figure 4A), no correlations were observed between IGF-I levels and changes in LDL or HDL cholesterol or Lp(a) concentrations ($r^2 < 0.10$ in all cases).

The plasma level of the bile acid intermediate, 7 α -hydroxy-4-cholesten-3-one (C4), reflects the activity of the rate-limiting enzyme in bile acid production, cholesterol 7 α -hydroxylase.⁴⁵ As seen in Table 1, GH treatment did not influence C4 levels in any group. The plasma ratio of unesterified lathosterol to cholesterol is a marker of hepatic and total body cholesterol synthesis.^{46,47} As displayed in Table 1, this marker was not influenced by GH treatment in any of the groups.

We finally investigated the effects of stepwise addition of GH, colestipol, and atorvastatin treatment in the girl with homozygous FH (Figure 5). Overall, there were no major

plasma lipoprotein changes. LDL cholesterol was reduced by 9% when all drugs were combined. Plasma triglycerides increased in response to GH, and this was normalized on combination with atorvastatin.

Discussion

GH exerts many important effects in the regulation of lipid metabolism.^{24,48} In general, GH increases the flux of energy in the lipid transport system by stimulating lipolysis in adipose tissue, fatty acid assimilation and triglyceride production in the liver, very-low-density lipoprotein secretion and catabolism, LDL clearance, and cholesterol excretion. Some of these effects are only observed in GH deficiency where they can be normalized by GH substitution, whereas others also occur in response to supraphysiological GH doses. The present study allows some important conclusions regarding the effects of GH on lipoprotein metabolism in humans.

First, administration of exogenous GH to healthy subjects resulted in a clear increase in the catabolism of LDL particles. We have reported that GH treatment for 5 days increases LDLR as measured directly in the liver of patients undergoing cholecystectomy¹² and that GH induces LDLR activity in cultured human hepatoma cells.⁴⁹ In the present experiment, the whole-body capacity for LDL clearance was examined through the LDL turnover technique, which reflects the total LDLR activity. The present data thus confirm and extend our earlier in vitro findings in human liver tissue. The FCR of LDL was increased by 27% in the normal subjects during GH treatment. This increase in FCR is of the same magnitude as the response seen with cholestyramine feeding at a dosage of 24 grams daily.⁸ However, the lowering of LDL cholesterol by GH was less than that observed in response to cholestyramine, probably because the production rate is somewhat increased with GH treatment. Stimulation of very-low-density lipoprotein production has been observed in response to GH substitution in adult GH-deficient patients using stable isotope kinetics.^{50,51} In our patient with pituitary insufficiency, LDL apoB FCR increased by 38% on GH substitution, despite the fact that he was administered a lower dose of GH than were the healthy subjects. In this single subject, LDL apoB synthesis was increased by GH, indicating that the flux of lipoproteins through plasma may be stimulated. Thus, a stimulation of LDL clearance by GH could be demonstrated, supporting the concept that GH is an important hormone in the complex physiological control of plasma LDL flux in humans.²⁴

Second, GH treatment resulted in a dose-related lowering of LDL cholesterol in younger and elderly normal male subjects. Although the response to the lowest dose was somewhat less pronounced in the elderly, this probably reflects a lower sensitivity to GH, as judged from the IGF-I response. The increase of LDL clearance that we demonstrate in young males can thus probably also be elicited in elderly males when the initial LDL clearance is reduced^{20,22} and when a relative deficiency of, or insensitivity to, GH is present.^{14,19} However, in both groups the LDL cholesterol lowering was relatively moderate ($\approx 15\%$ at the highest dose). The LDLR stimulatory effect seen with GH treatment may have several explanations. Hypothetically, this effect could be mediated by

Effects of GH on Plasma Lipids, Lipoproteins, and Sterol Synthesis Markers

	Group	Basal	Week 1	Week 2	Week 3
Cholesterol (mmol/L)	Younger	4.8±0.3	4.5±0.3	4.6±0.3	4.3±0.4*
	Elderly	5.7±0.4	5.6±0.3	5.5±0.3	5.2±0.3*
	FH patients	10.5±0.6	10.3±0.6	10.2±0.6	9.5±0.4*
LDL cholesterol (mmol/L)	Younger	3.2±0.2	2.9±0.2	2.9±0.2	2.6±0.2**
	Elderly	3.9±0.3	3.9±0.3	3.7±0.2	3.4±0.3**
	FH patients	8.7±0.6	8.3±0.5	8.2±0.6	7.4±0.4**
HDL cholesterol (mmol/L)	Younger	1.3±0.1	1.2±0.1**	1.1±0.1**	1.1±0.1**
	Elderly	1.5±0.1	1.3±0.1**	1.2±0.1**	1.3±0.1**
	FH patients	1.3±0.1	1.2±0.1**	1.2±0.1**	1.1±0.1**
Triglycerides (mmol/L)	Younger	1.0±0.1	1.0±0.1	1.2±0.2***	1.5±0.3***
	Elderly	0.9±0.1	1.2±0.1*	1.5±0.1***	1.6±0.2***
	FH patients	1.6±0.2	1.9±0.2*	2.0±0.4***	2.5±0.5***
apoAI (g/L)	Younger	1.24±0.1	1.20±0.1	1.26±0.1	1.20±0.1
	Elderly	1.25±0.1	1.18±0.1	1.22±0.1	1.19±0.1**
	FH patients	1.13±0.1	1.20±0.1*	1.16±0.1	1.14±0.1
apoB (g/L)	Younger	0.81±0.1	0.79±0.1	0.79±0.04	0.73±0.1***
	Elderly	1.03±0.1	0.99±0.04	0.98±0.1	0.89±0.1***
	FH patients	2.05±0.2	2.04±0.1	1.91±0.1	1.89±0.1***
Lp(a) (mg/L)	Younger	374 (53–630)	414 (53–691)**	449 (41–831)***	449 (126–767)***
	Elderly	66 (30–565)	91 (30–619)**	149 (30–664)***	182 (30–759)***
	FH patients	376 (30–1032)	410 (30–1218)**	474 (30–1230)***	474 (30–1297)***
C4 (ng/mL)	Younger	12±3	14±4	9±1	9±3
	Elderly	12±3	16±4	10±3	11±4
	FH patients	18±6	18±3	14±2	14±4
C4/cholesterol (ng/mg)	Younger	6.2±1.4	8.0±2.0	5.0±0.5	5.3±1.6
	Elderly	5.7±1.3	7.3±2.0	4.8±1.5	5.1±2.2
	FH patients	4.6±1.7	4.5±0.7	3.5±0.6	3.9±1.2
Iatho/cholesterol (ng/mg)	Younger	560±79	540±59	530±41	550±79
	Elderly	590±44	530±36	590±89	440±85
	FH patients	670±88	590±79	560±67	620±80

All data are expressed as mean±SEM except for Lp(a), which is presented as median (range).

C4 indicates 7 α OH-4-cholesten-3-one.

P-values significantly different from basal are: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

IGF-I, a somatomedin that increases in plasma after GH secretion. In the present study, we were able to demonstrate a 3-fold increase of IGF-I in response to the GH treatment, but there was no correlation between IGF-I levels and lipoprotein changes. Recent studies in humans⁵² and in rats⁵³ have demonstrated that IGF-I treatment does not result in the same effects as GH treatment on plasma lipoprotein levels or on hepatic LDLR expression in vivo, which strongly indicates that the effect of GH treatment on LDLR and LDL catabolism is direct. The molecular mechanisms for this effect remain to be explored.

Third, the effects of GH on LDL cholesterol in heterozygous FH were similar to those seen in younger and elderly healthy subjects. Reductions in LDL cholesterol have also been observed in FH heterozygotes treated with 0.05 IU/kg

per day for 12 weeks.⁵⁴ Although probably of limited therapeutic value in heterozygous FH, it is evident that the capacity for LDL-lowering by GH was not decreased, despite a reduced initial expression of LDLR. However, our data from the receptor-deficient FH homozygote clearly indicate that the capability to express LDLR is a prerequisite for LDL-lowering effects of GH. Although only 1 patient was studied, this finding is in clear contrast to observations in LDLR-deficient (knockout) mice, in which GH treatment reduced plasma LDL cholesterol levels.²⁶ Thus, there is a clear species difference in the lipid-lowering effect of GH. The explanation to this is unclear, but it may be mentioned that in rodents GH stimulates the hepatic editing of apoB, resulting in a faster clearance of hepatic lipoproteins from the circulation.^{53,55} This response should not occur in humans.⁵⁶




Figure 5. The levels of plasma LDL and HDL cholesterol and triglycerides in the child with homozygous FH during the study. Treatment regimens are indicated. Atorvastatin was administered at a dosage of 0.6 mg/kg per day. GH was administered at the indicated doses (IU/kg per day). Colestipol was successively increased to 0.45 g/kg per day during the first 10 days. No treatment was administered during the washout period.

Also, the different effect of GH in stimulation of bile acid synthesis may be of importance for this species difference.

Fourth, the administration of GH to normal human subjects does not increase bile acid synthesis, which is in agreement with previous results in which bile acid production was studied using isotope kinetics in young normal male subjects before and during treatment with GH.¹⁵ In the rat, bile acid synthesis is dependent on the presence of GH,²⁵ and stimulation of cholesterol breakdown to bile acids can be achieved by pharmacological doses of GH.²³ Thus, there is a difference between rodents and humans regarding the regulation of bile acid synthesis by GH. The metabolic basis behind this is unclear, but it is of interest to note that species differences have recently been described in the molecular regulation of bile acid synthesis.^{57–61} Furthermore, there appear to be species differences in the plasma lipoprotein response to stimulation of bile acid synthesis in LDLR deficiency.⁶² Thus, stimulation of cholesterol 7 α -hydroxylase by gene transfer,⁶³ GH treatment,²⁶ or interruption of bile acid intestinal uptake⁴⁵ lowers LDL cholesterol in LDLR-deficient knockout mice, whereas complete biliary diversion in homozygous FH does not lower plasma LDL.⁶⁴

Fifth, the concentration of plasma Lp(a) increased dose-dependently in all groups. Also in GH deficiency, Lp(a) levels are increased by GH substitution,^{17,65–67} whereas they are reduced after therapy of acromegaly.^{68,69} It has been shown that IGF-I does not increase Lp(a). Instead, its administration reduces Lp(a).^{52,70–72} Therefore, the present results strongly support the concept that GH directly stimulates the synthesis and secretion of apo(a), a mechanism demonstrated in mice transgenic for human apo(a).⁷³ Although the clinical relevance of the GH-induced Lp(a) increase is not known, this consistent effect together with increased plasma levels of glucose, triglycerides, and reduced plasma HDL cholesterol should call for some caution in situations in which GH treatment is considered.

Acknowledgments

The skilful technical assistance of Lilian Larsson, Danuta Cosliani, and Sabine Süllo-Barin, and the manuscript preparation of Lena Emtestam are gratefully acknowledged. The studies were supported by grants from the Swedish Medical Research Council (03X-7137), the Swedish Heart–Lung Foundation, the Loo and Hans Osterman

and the Thuring Foundations, the Foundation of Old Female Servants, and the Karolinska Institute. Human growth hormone was kindly provided by Pharmacia, Stockholm, Sweden.

References

- Goldstein JL, Hobbs HH, Brown MS. Familial hypercholesterolemia. In: Scriver CR, Beaudet AL, Sly WS, Valle D, eds. *The Metabolic and Molecular Basis of Inherited Disease*. New York: McGraw-Hill; 2001: 2863–2913.
- Myant NB. *Cholesterol Metabolism, LDL, and the LDL Receptor*. San Diego: Academic Press; 1990.
- Brown MS, Goldstein JL. A receptor-mediated pathway for cholesterol homeostasis. *Science*. 1986;232:34–47.
- Reihner E, Rudling M, Ståhlberg D, Berglund L, Ewerth S, Björkhem I, Einarsson K, Angelin B. Influence of pravastatin, a specific inhibitor of HMG-CoA reductase, on hepatic metabolism of cholesterol. *N Engl J Med*. 1990;323:224–228.
- Rudling M, Reihner E, Einarsson K, Ewerth S, Angelin B. Low density lipoprotein receptor-binding activity in human tissues: Quantitative importance of hepatic receptors and evidence for regulation of their expression in vivo. *Proc Natl Acad Sci U S A*. 1990;87:3469–3473.
- Rudling M, Angelin B, Ståhle L, Reihner E, Sahlin S, Olivecrona H, Björkhem I, Einarsson C. Regulation of hepatic low-density lipoprotein receptor, 3-hydroxy-3-methylglutaryl coenzyme A reductase, and cholesterol 7 α -hydroxylase mRNAs in human liver. *J Clin Endocrinol Metab*. 2002;87:4307–4313.
- Angelin B, Olivecrona H, Reihner E, Rudling M, Ståhlberg D, Eriksson M, Ewerth S, Henriksson P, Einarsson K. Hepatic cholesterol metabolism in estrogen-treated men. *Gastroenterology*. 1992;103:1657–1663.
- Shepherd J, Packard CJ, Bicker S, Lawrie TD, Morgan HG. Cholestyramine promotes receptor-mediated low-density-lipoprotein catabolism. *N Engl J Med*. 1980;302:1219–1222.
- Bilheimer DW, Grundy SM, Brown MS, Goldstein JL. Mevinolin and colestipol stimulate receptor-mediated clearance of low density lipoprotein from plasma in familial hypercholesterolemia heterozygotes. *Proc Natl Acad Sci U S A*. 1983;80:4124–4128.
- Eriksson M, Berglund L, Rudling M, Henriksson P, Angelin B. Effects of estrogen on low density lipoprotein metabolism in males. Short-term and long-term studies during hormonal treatment of prostatic carcinoma. *J Clin Invest*. 1989;84:802–810.
- Walsh BW, Schiff I, Rosner B, Greenberg L, Ravnkar V, Sacks FM. Effects of postmenopausal estrogen replacement on the concentrations and metabolism of plasma lipoproteins. *N Engl J Med*. 1991;325: 1196–1204.
- Rudling M, Norstedt G, Olivecrona H, Reihner E, Gustafsson JA, Angelin B. Importance of growth hormone for the induction of hepatic low density lipoprotein receptors. *Proc Natl Acad Sci U S A*. 1992;89:6983–6987.
- Friedman M, Byers SO, Rosenman RH, Li CH, Neuman R. Effect of subacute administration of human growth hormone on various serum lipid and hormone levels of hypercholesterolemic and normocholesterolemic subjects. *Metabolism*. 1974;23:905–912.
- Rudman D, Kutner MH, Rogers CM, Lubin MF, Fleming GA, Bain RP. Impaired growth hormone secretion in the adult population: relation to age and adiposity. *J Clin Invest*. 1981;67:1361–1369.
- Olivecrona H, Ericsson S, Angelin B. Growth hormone treatment does not alter biliary lipid metabolism in healthy adult men. *J Clin Endocrinol Metab*. 1995;80:1113–1117.
- Salomon F, Cuneo RC, Hesp R, Sonksen PH. The effects of treatment with recombinant human growth hormone on body composition and metabolism in adults with growth hormone deficiency. *N Engl J Med*. 1989;321:1797–1803.
- Eden S, Wiklund O, Oscarsson J, Rosen T, Bengtsson BA. Growth hormone treatment of growth hormone-deficient adults results in a marked increase in Lp(a) and HDL cholesterol concentrations. *Arterioscler Thromb*. 1993;13:296–301.
- Russell-Jones DL, Watts GF, Weissberger A, Naoumova R, Myers J, Thompson GR, Sonksen PH. The effect of growth hormone replacement on serum lipids, lipoproteins, apolipoproteins and cholesterol precursors in adult growth hormone deficient patients. *Clin Endocrinol (Oxf)*. 1994; 41:345–350.
- Vahl N, Klausen I, Christiansen JS, Jorgensen JO. Growth hormone (GH) status is an independent determinant of serum levels of cholesterol and triglycerides in healthy adults. *Clin Endocrinol (Oxf)*. 1999;51:309–316.

20. Ericsson S, Eriksson M, Vitols S, Einarsson K, Berglund L, Angelin B. Influence of age on the metabolism of plasma low density lipoproteins in healthy males. *J Clin Invest.* 1991;87:591–596.
21. Einarsson K, Nilsell K, Leijb B, Angelin B. Influence of age on secretion of cholesterol and synthesis of bile acids by the liver. *N Engl J Med.* 1985;313:277–282.
22. Grundy SM, Vega GL, Bilheimer DW. Kinetic mechanisms determining variability in low density lipoprotein levels and rise with age. *Arteriosclerosis.* 1985;5:623–630.
23. Parini P, Angelin B, Rudling M. Cholesterol and lipoprotein metabolism in aging: reversal of hypercholesterolemia by growth hormone treatment in old rats. *Arterioscler Thromb Vasc Biol.* 1999;19:832–839.
24. Angelin B, Rudling M. Growth hormone and hepatic lipoprotein metabolism. *Curr Opin Lipidol.* 1994;5:160–165.
25. Rudling M, Parini P, Angelin B. Growth hormone and bile acid synthesis. Key role for the activity of hepatic microsomal cholesterol 7 α -hydroxylase in the rat. *J Clin Invest.* 1997;99:2239–2245.
26. Rudling M, Angelin B. Growth hormone reduces plasma cholesterol in LDL receptor-deficient mice. *FASEB J.* 2001;15:1350–1356.
27. Wiklund O, Angelin B, Olofsson SO, Eriksson M, Fager G, Berglund L, Bondjers G. Apolipoprotein(a) and ischaemic heart disease in familial hypercholesterolaemia. *Lancet.* 1990;335:1360–1363.
28. Lind S, Eriksson M, Rystedt E, Wiklund O, Angelin B, Eggertsen G. Low frequency of the common Norwegian and Finnish LDL-receptor mutations in Swedish patients with familial hypercholesterolaemia. *J Intern Med.* 1998;244:19–25.
29. Lind S, Rystedt E, Eriksson M, Wiklund O, Angelin B, Eggertsen G. Genetic characterization of Swedish patients with familial hypercholesterolemia: a heterogeneous pattern of mutations in the LDL receptor gene. *Atherosclerosis.* 2002;163:399–407.
30. Gutierrez G, Schneider A, Jobs J, Schmidt H, Korte A, Manns MP, Stuhmann M. Homozygous familial hypercholesterolemia: A novel point mutation (W556R) in a Turkish patient. *Hum Mutat.* 2000;16:374.
31. Bang P, Eriksson U, Sara V, Wivall IL, Hall K. Comparison of acid ethanol extraction and acid gel filtration prior to IGF-I and IGF-II radioimmunoassays: improvement of determinations in acid ethanol extracts by the use of truncated IGF-I as radioligand. *Acta Endocrinol (Copenh).* 1991;124:620–629.
32. Pova G, Roovete A, Hall K. Cross-reaction of serum somatomedin-binding protein in a radioimmunoassay developed for somatomedin-binding protein isolated from human amniotic fluid. *Acta Endocrinol (Copenh).* 1984;107:563–570.
33. Carlson K. Lipoprotein fractionation. *J Clin Pathol Suppl (Assoc Clin Pathol).* 1973;5:32–37.
34. Lopes-Virella MF, Stone P, Ellis S, Colwell JA. Cholesterol determination in high-density lipoproteins separated by three different methods. *Clin Chem.* 1977;23:882–884.
35. Galman C, Arvidsson I, Angelin B, Rudling M. Monitoring hepatic cholesterol 7 α -hydroxylase activity by assay of the stable bile acid intermediate 7 α -hydroxy-4-cholesten-3-one in peripheral blood. *J Lipid Res.* 2003;44:859–866.
36. Lund E, Sisfontes L, Reihner E, Bjorkhem I. Determination of serum levels of unesterified lathosterol by isotope dilution-mass spectrometry. *Scand J Clin Lab Invest.* 1989;49:165–171.
37. Bilheimer DW, Eisenberg S, Levy RI. The metabolism of very low density lipoprotein proteins. I. Preliminary in vitro and in vivo observations. *Biochim Biophys Acta.* 1972;260:212–221.
38. Matthews CME. The theory of tracer experiments with 131-labeled plasma proteins. *Phys Med Biol.* 1957;2:36–53.
39. Langer T, Strober W, Levy RI. The metabolism of low density lipoprotein in familial type II hyperlipoproteinemia. *J Clin Invest.* 1972;51:1528–1536.
40. Kesaniemi YA, Grundy SM. Significance of low density lipoprotein production in the regulations of plasma cholesterol level in man. *J Clin Invest.* 1982;70:13–21.
41. Juul A, Main K, Blum WF, Lindholm J, Ranke MB, Skakkebaek NE. The ratio between serum levels of insulin-like growth factor (IGF)-I and the IGF binding proteins (IGFBP-1, 2 and 3) decreases with age in healthy adults and is increased in acromegalic patients. *Clin Endocrinol (Oxf).* 1994;41:85–93.
42. Berglund L, Witztum JL, Galeano NF, Khouw AS, Ginsberg HN, Ramakrishnan R. Three-fold effect of lovastatin treatment on low density lipoprotein metabolism in subjects with hyperlipidemia: increase in receptor activity, decrease in apoB production, and decrease in particle affinity for the receptor. Results from a novel triple-tracer approach. *J Lipid Res.* 1998;39:913–924.
43. Witztum JL, Young SG, Elam RL, Carew TE, Fisher M. Cholestyramine-induced changes in low density lipoprotein composition and metabolism. I. Studies in the guinea pig. *J Lipid Res.* 1985;26:92–103.
44. Ericsson S, Berglund L, Frostegard J, Einarsson K, Angelin B. The influence of age on low density lipoprotein metabolism: effects of cholestyramine treatment in young and old healthy male subjects. *J Intern Med.* 1997;242:329–337.
45. Galman C, Ostlund-Lindqvist AM, Bjorquist A, Schreyer S, Svensson L, Angelin B, Rudling M. Pharmacological interference with intestinal bile acid transport reduces plasma cholesterol in LDL receptor/apoE deficiency. *FASEB J.* 2003;17:265–267.
46. Bjorkhem I, Miettinen T, Reihner E, Ewerth S, Angelin B, Einarsson K. Correlation between serum levels of some cholesterol precursors and activity of HMG-CoA reductase in human liver. *J Lipid Res.* 1987;28:1137–1143.
47. Kempen HJ, Glatz JF, Gevers Leuven JA, van der Voort HA, Katan MB. Serum lathosterol concentration is an indicator of whole-body cholesterol synthesis in humans. *J Lipid Res.* 1988;29:1149–1155.
48. Corpas E, Harman SM, Blackman MR. Human growth hormone and human aging. *Endocrinol Rev.* 1993;14:20–39.
49. Parini P, Angelin B, Lobie PE, Norstedt G, Rudling M. Growth hormone specifically stimulates the expression of low density lipoprotein receptors in human hepatoma cells. *Endocrinology.* 1995;136:3767–3773.
50. Christ ER, Cummings MH, Albany E, Uempleby AM, Lumb PJ, Wierzbicki AS, Naoumova RP, Boroujerdi MA, Sonksen PH, Russell-Jones DL. Effects of growth hormone (GH) replacement therapy on very low density lipoprotein apolipoprotein B100 kinetics in patients with adult GH deficiency: a stable isotope study. *J Clin Endocrinol Metab.* 1999;84:307–316.
51. Kearney T, de Gallegos CN, Proudler A, Parker K, Anayaoku V, Bannister P, Venkatesan S, Johnston DG. Effects of short- and long-term growth hormone replacement on lipoprotein composition and on very-low-density lipoprotein and low-density lipoprotein apolipoprotein B100 kinetics in growth hormone-deficient hypopituitary subjects. *Metabolism.* 2003;52:50–59.
52. Olivecrona H, Johansson AG, Lindh E, Ljunghall S, Berglund L, Angelin B. Hormonal regulation of serum lipoprotein(a) levels. Contrasting effects of growth hormone and insulin-like growth factor-I. *Arterioscler Thromb Vasc Biol.* 1995;15:847–849.
53. Rudling M, Olivecrona H, Eggertsen G, Angelin B. Regulation of rat hepatic low density lipoprotein receptors. In vivo stimulation by growth hormone is not mediated by insulin-like growth factor I. *J Clin Invest.* 1996;97:292–299.
54. Tonstad S, Sundt E, Ose L, Hagve TA, Fruchart JC, Bard JM, Eden S. The effect of growth hormone on low-density lipoprotein cholesterol and lipoprotein (a) levels in familial hypercholesterolemia. *Metabolism.* 1996;45:1415–1421.
55. Sjoberg A, Oscarsson J, Bostrom K, Innerarity TL, Eden S, Olofsson SO. Effects of growth hormone on apolipoprotein-B (apoB) messenger ribonucleic acid editing, and apoB 48 and apoB 100 synthesis and secretion in the rat liver. *Endocrinology.* 1992;130:3356–3364.
56. Davidson NO. Apolipoprotein B mRNA editing: a key controlling element targeting fats to proper tissue. *Ann Med.* 1993;25:539–543.
57. Bjorkhem I, Araya Z, Rudling M, Angelin B, Einarsson C, Wikvall K. Differences in the regulation of the classical and the alternative pathway for bile acid synthesis in human liver. No coordinate regulation of CYP7A1 and CYP27A1. *J Biol Chem.* 2002;277:26804–26807.
58. Sinal CJ, Yoon M, Gonzalez FJ. Antagonism of the actions of peroxisome proliferator-activated receptor- α by bile acids. *J Biol Chem.* 2001;276:47154–47162.
59. Pineda Torra I, Claudel T, Duval C, Kosykh V, Fruchart JC, Staels B. Bile Acids Induce the Expression of the Human Peroxisome Proliferator-Activated Receptor α Gene via Activation of the Farnesoid X Receptor. *Mol Endocrinol.* 2003;17:259–272.
60. Chiang JY, Kimmel R, Stroup D. Regulation of cholesterol 7 α -hydroxylase gene (CYP7A1) transcription by the liver orphan receptor (LXR α). *Gene.* 2001;262:257–265.
61. Peet DJ, Turley SD, Ma W, Janowski BA, Lobaccaro JM, Hammer RE, Mangelsdorf DJ. Cholesterol and bile acid metabolism are impaired in mice lacking the nuclear oxysterol receptor LXR α . *Cell.* 1998;93:693–704.
62. Angelin B, Eriksson M, Rudling M. Bile acids and lipoprotein metabolism: a renaissance for bile acids in the post-statin era? *Curr Opin Lipidol.* 1999;10:269–274.
63. Spady DK, Cuthbert JA, Willard MN, Meidell RS. Overexpression of cholesterol 7 α -hydroxylase (CYP7A) in mice lacking the low density

- lipoprotein (LDL) receptor gene. LDL transport and plasma LDL concentrations are reduced. *J Biol Chem.* 1998;273:126–132.
64. Deckelbaum RJ, Lees RS, Small DM, Hedberg SE, Grundy SM. Failure of complete bile diversion and oral bile acid therapy in the treatment of homozygous familial hypercholesterolemia. *N Engl J Med.* 1977;296:465–470.
 65. O'Halloran DJ, Wieringa G, Tsatsoulis A, Shalet SM. Increased serum lipoprotein(a) concentrations after growth hormone (GH) treatment in patients with isolated GH deficiency. *Ann Clin Biochem.* 1996;33:330–334.
 66. Johannsson G, Oscarsson J, Rosen T, Wiklund O, Olsson G, Wilhelmssen L, Bengtsson BA. Effects of 1 year of growth hormone therapy on serum lipoprotein levels in growth hormone-deficient adults. Influence of gender and Apo(a) and ApoE phenotypes. *Arterioscler Thromb Vasc Biol.* 1995;15:2142–2150.
 67. Garry P, Collins P, Devlin JG. An open 36-month study of lipid changes with growth hormone in adults: lipid changes following replacement of growth hormone in adult acquired growth hormone deficiency. *Eur J Endocrinol.* 1996;134:61–66.
 68. Lam KS, Pang RW, Janus ED, Kung AW, Wang CC. Serum apolipoprotein(a) correlates with growth hormone levels in Chinese patients with acromegaly. *Atherosclerosis.* 1993;104:183–188.
 69. Wildbrett J, Hanefeld M, Fucker K, Pinzer T, Bergmann S, Siegert G, Breidert M. Anomalies of lipoprotein pattern and fibrinolysis in acromegalic patients: relation to growth hormone levels and insulin-like growth factor I. *Exp Clin Endocrinol Diabetes.* 1997;105:331–335.
 70. Oscarsson J, Lundstam U, Gustafsson B, Wilton P, Eden S, Wiklund O. Recombinant human insulin-like growth factor-I decreases serum lipoprotein(a) concentrations in normal adult men. *Clin Endocrinol (Oxf).* 1995;42:673–676.
 71. Laron Z, Wang XL, Klinger B, Silbergeld A, Wilcken DE. Insulin-like growth factor-I decreases serum lipoprotein (a) during long-term treatment of patients with Laron syndrome. *Metabolism.* 1996;45:1263–1266.
 72. Laron Z, Wang XL, Klinger B, Silbergeld A, Wilcken DE. Growth hormone increases and insulin-like growth factor-I decreases circulating lipoprotein(a). *Eur J Endocrinol.* 1997;136:377–381.
 73. Tao R, Acquati F, Marcovina SM, Hobbs HH. Human growth hormone increases apo(a) expression in transgenic mice. *Arterioscler Thromb Vasc Biol.* 1999;19:2439–2447.

Arteriosclerosis, Thrombosis, and Vascular Biology



JOURNAL OF THE AMERICAN HEART ASSOCIATION

Growth Hormone Induces Low-Density Lipoprotein Clearance but not Bile Acid Synthesis in Humans

Suzanne Lind, Mats Rudling, Sverker Ericsson, Hans Olivecrona, Mats Eriksson, Birgit Borgström, Gösta Eggertsen, Lars Berglund and Bo Angelin

Arterioscler Thromb Vasc Biol. 2004;24:349-356; originally published online December 4, 2003;

doi: 10.1161/01.ATV.0000110657.67317.90

Arteriosclerosis, Thrombosis, and Vascular Biology is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2003 American Heart Association, Inc. All rights reserved.

Print ISSN: 1079-5642. Online ISSN: 1524-4636

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://atvb.ahajournals.org/content/24/2/349>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Arteriosclerosis, Thrombosis, and Vascular Biology* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:

<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Arteriosclerosis, Thrombosis, and Vascular Biology* is online at:

<http://atvb.ahajournals.org/subscriptions/>