

Protease-Activated Receptor-1 Signaling The Big Picture

Robert Flaumenhaft

Proteases serve an essential role in blood clotting, inflammation, angiogenesis, wound healing, and other vascular processes. The endothelium must be able to sense and respond to vascular proteases to coordinate these activities. Protease-activated receptors (PARs) are G-protein-coupled receptors responsible for sensing proteases, and PAR1 is a critical family member on endothelium. Like other PARs, endothelial PAR1 is able to sense the proteolytic environment by virtue of an unusual feature: an N terminus that serves as a substrate for many different proteases. Cleavage of the N terminus reveals a tethered ligand capable of activating the receptor via an intramolecular signaling mechanism (Figure).¹ More recent studies have shown that different proteases have unique cleavage sites, leading to distinct tethered ligands. Cleavage of endothelial PAR1 by thrombin results in apoptosis, release of granules, surface expression of inflammatory markers, and loss of barrier function.³ In contrast, cleavage of PAR1 by activated protein C (APC) can result in protection from apoptotic stimuli, suppression of inflammatory markers, and fortification of barrier function.³ These observations raise the complex question of how endothelial cells integrate PAR1 stimulation to mediate such varied responses.

See accompanying article on page 1891

Strategies to define the intricate signaling pathways downstream of PAR1 have evolved over the years. Initial efforts defined the specific G proteins that couple to PAR1. Consistent with its ability to direct a broad range of cellular responses, PAR1 was found to couple with several G proteins, including $G\alpha_q$, $G\alpha_{12/13}$, and $G\alpha_i$, as well as $G\beta\gamma$.³ Later, PAR1 coupling to β -arrestins was demonstrated in the context of APC stimulation.⁴ Many downstream phosphorylation events have been characterized primarily via the use of phospho-specific antibodies. Although this approach provides useful information regarding the kinetics and dose-dependency of endothelial PAR1 signaling, it focuses on only a small component of the signaling pathway without providing any sense of how the components interact. As a result, this approach cannot comprehensively compare signaling pathways induced by different PAR1 agonists nor can it comprehensively assess the effect of different PAR1 antagonists on downstream signaling.

van den Biggelaar et al^{2,5} at Sanquin Research in Amsterdam have now applied a powerful global phosphoproteomic approach to evaluate signaling cascades activated after PAR1 stimulation in endothelial cells. To enable quantitative analysis of phosphorylated proteins, the authors used a 3-way reverse SILAC (stable isotope labeling with amino acids in cell culture) labeling strategy requiring metabolic labeling of blood outgrowth endothelial cells with light, medium, or heavy arginine and lysine, followed by phosphopeptide separation and analysis using high-resolution mass spectrometry. This rigorous approach enabled stringent criteria for quantifying phosphorylation events. The authors quantified 2553 phosphosites altogether and, importantly, 268 proteins (410 phosphosites) that demonstrated time-dependent changes (either increased or decreased) in phosphorylation after thrombin stimulation.²

The authors applied this phosphoproteomics strategy to address several issues in the PAR1 signaling field. First, they focused on an assertion arising from prior reports that thrombin modulates downstream signaling differently than SFLLRN (serine-phenylalanine-leucine-leucine-arginine-asparagine), a PAR1-activating peptide based on the tethered ligand produced by thrombin cleavage. A previous study showed that signaling through $G\alpha_{12/13}$ required lower concentrations of thrombin than signaling through $G\alpha_q$, whereas SFLLRN-induced signaling showed the opposite pattern.⁶ In contrast, the author's proteomic studies showed essentially identical phosphoprotein profiles after exposure to thrombin versus SFLLRN. One possible explanation for the apparent discrepancy is that the proteomic study was performed at near saturating concentrations of agonists. Nonetheless, the results provide investigators in the field assurance that this PAR1-activating peptide can be used as a reliable mimetic of thrombin with regard to PAR1 signaling function.

The authors also used the phosphoproteomic approach to compare and contrast PAR1 antagonists. The authors compared vorapaxar (an orthosteric inhibitor that associates with the shallow ligand-binding pocket of PAR1)⁷ with parmodulin-2 (an antagonist that associates with the cytosolic face of PAR1 and blocks some, but not all, downstream pathways).⁸⁻¹⁰ This analysis demonstrated differential inhibition of thrombin-mediated stimulation by parmodulin-2 compared with vorapaxar. Vorapaxar inhibited essentially all downstream signaling, buttressing the contention that PAR1 mediates the vast majority of thrombin signaling in endothelium. Parmodulin-2 blocked many PAR1-mediated phosphorylation events but was less effective at blocking dephosphorylation events that occurred in response to thrombin. This unexpected observation indicates that parmodulin-2 does not block many thrombin-activated phosphatase activities, providing additional complexity to the previously reported inhibitory bias of

From the Beth Israel Deaconess Medical Center, Department of Medicine, Harvard Medical School, Boston, MA.

Correspondence to Robert Flaumenhaft, Beth Israel Deaconess Medical Center, Department of Medicine, 330 Brookline Ave, Boston, MA 02215. E-mail rflaumen@bidmc.harvard.edu

Arterioscler Thromb Vasc Biol. 2017;37:1809-1811.
DOI: 10.1161/ATVBAHA.117.310068.)

© 2017 American Heart Association, Inc.

Arterioscler Thromb Vasc Biol is available at <http://atvb.ahajournals.org>
DOI: 10.1161/ATVBAHA.117.310068




Figure. The N-terminus of protease-activated receptor (PAR1; red) is cleaved revealing a tethered ligand (yellow), which interacts with the ligand-binding site. Ligation of this binding site results in a conformational change in the receptor that signals to G-proteins and β -arrestin stimulating an intricate network of phosphorylation events (see van den Eshof et al² for details). PAR1 signaling coordinates several endothelial cell functions.

parmodulin-2.¹⁰ These results underscore the value of global phosphoproteomics in assessing the mechanism of novel antagonists.

Another important area in PAR1 biology is biased signaling. For example, in explaining why cleavage of PAR1 by thrombin results in different endothelial responses than those observed after APC cleavage, some investigators have shown that thrombin-mediated signaling is biased toward G proteins, and APC-mediated signaling is biased toward β -arrestins.^{3,11} APC, matrix metalloproteinase 1, proteinase 3, and neutrophil elastase all cleave PAR1 at distinct sites.² Peptides based on these noncanonical tethered ligands stimulate PAR1. When the authors compared the different PAR1-stimulating peptides to SFLLRN, they found that changes in the phosphoproteome induced by exposure to noncanonical peptides were minor compared with the robust changes induced by SFLLRN, and changes were generally in the same direction for both SFLLRN and noncanonical ligands. Surprisingly, only a few phosphosites were found that diverged from this pattern. This result is inconsistent with previous work using phosphorylation-specific antibodies that showed distinct phosphorylation events after exposure to noncanonical ligands compared with those after exposure to SFLLRN.

So how does one square previously reported results with these phosphoproteomic data? One possibility is timing. Because noncanonical peptide ligands can take longer to stimulate signaling, their distinct phosphorylation events may not have been captured. Another possibility is sensitivity (ie, phospho-specific antibodies may be more sensitive). Differences in cell type and origin could also contribute to the differences with previously published results. Finally, it is important to remember that not all signaling events are phosphorylation events. Differences in functional responses could be related to alternative second messenger systems. Nonetheless, these phosphoproteomic data warrant a critical re-evaluation of the nature of noncanonical PAR1 signal transduction in endothelial cells. Indeed, this rigorous phosphoproteomic approach provides investigators with a whole new way to look at PAR1 signaling—the big picture.

Acknowledgments

R. Flaumenhaft is supported by R35 HL135775 and R01 HL125275.

References

1. Vu TK, Hung DT, Wheaton VI, Coughlin SR. Molecular cloning of a functional thrombin receptor reveals a novel proteolytic mechanism of receptor activation. *Cell*. 1991;64:1057–1068.
2. van den Eshof BL, Hoogendijk AJ, Simpson PJ, van Alphen FPJ, Zanivan S, Mertens K, Meijer AB, van den Biggelaar M. Paradigm of biased PAR1 (protease-activated receptor-1) activation and inhibition in endothelial cells dissected by phosphoproteomics. *Arterioscler Thromb Vasc Biol*. 2017;37:1891–1902. doi: 10.1161/ATVBAHA.117.309926.
3. Flaumenhaft R, De Ceunynck K. Targeting PAR1: now what? *Trends Pharmacol Sci*. 2017;38:701–716. doi: 10.1016/j.tips.2017.05.001.
4. Soh UJ, Trejo J. Activated protein C promotes protease-activated receptor-1 cytoprotective signaling through β -arrestin and dishevelled-2 scaffolds. *Proc Natl Acad Sci USA*. 2011;108:E1372–E1380. doi: 10.1073/pnas.1112482108.
5. van den Biggelaar M, Hernández-Fernaud JR, van den Eshof BL, Neilson LJ, Meijer AB, Mertens K, Zanivan S. Quantitative phosphoproteomics unveils temporal dynamics of thrombin signaling in human endothelial cells. *Blood*. 2014;123:e22–e36. doi: 10.1182/blood-2013-12-546036.
6. McLaughlin JN, Shen L, Holinstat M, Brooks JD, Dibenedetto E, Hamm HE. Functional selectivity of G protein signaling by agonist peptides and thrombin for the protease-activated receptor-1. *J Biol Chem*. 2005;280:25048–25059. doi: 10.1074/jbc.M414090200.
7. Zhang C, Srinivasan Y, Arlow DH, Fung JJ, Palmer D, Zheng Y, Green HF, Pandey A, Dror RO, Shaw DE, Weis WI, Coughlin SR, Kobilka BK. High-resolution crystal structure of human protease-activated receptor 1. *Nature*. 2012;492:387–392. doi: 10.1038/nature11701.
8. Dowal L, Sim DS, Dilks JR, Blair P, Beaudry S, Denker BM, Koukos G, Kuliopulos A, Flaumenhaft R. Identification of an antithrombotic allosteric modulator that acts through helix 8 of PAR1. *Proc Natl Acad Sci USA*. 2011;108:2951–2956. doi: 10.1073/pnas.1014863108.
9. Dockendorff C, Aisiku O, Verplank L, Dilks JR, Smith DA, Gunnink SF, Dowal L, Negri J, Palmer M, Macpherson L, Schreiber SL, Flaumenhaft R. Discovery of 1,3-diaminobenzenes as selective inhibitors of platelet activation at the PAR1 receptor. *ACS Med Chem Lett*. 2012;3:232–237. doi: 10.1021/ml2002696.
10. Aisiku O, Peters CG, De Ceunynck K, Ghosh CC, Dilks JR, Fustolo-Gunnink SF, Huang M, Dockendorff C, Parikh SM, Flaumenhaft R. Parmodulins inhibit thrombus formation without inducing endothelial injury caused by vorapaxar. *Blood*. 2015;125:1976–1985. doi: 10.1182/blood-2014-09-599910.
11. Mosnier LO, Sinha RK, Burnier L, Bouwens EA, Griffin JH. Biased agonism of protease-activated receptor 1 by activated protein C caused by noncanonical cleavage at Arg46. *Blood*. 2012;120:5237–5246. doi: 10.1182/blood-2012-08-452169.

Arteriosclerosis, Thrombosis, and Vascular Biology



JOURNAL OF THE AMERICAN HEART ASSOCIATION

Protease-Activated Receptor-1 Signaling: The Big Picture Robert Flaumenhaft

Arterioscler Thromb Vasc Biol. 2017;37:1809-1811

doi: 10.1161/ATVBAHA.117.310068

Arteriosclerosis, Thrombosis, and Vascular Biology is published by the American Heart Association, 7272
Greenville Avenue, Dallas, TX 75231

Copyright © 2017 American Heart Association, Inc. All rights reserved.

Print ISSN: 1079-5642. Online ISSN: 1524-4636

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://atvb.ahajournals.org/content/37/10/1809>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Arteriosclerosis, Thrombosis, and Vascular Biology* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Arteriosclerosis, Thrombosis, and Vascular Biology* is online at:
<http://atvb.ahajournals.org/subscriptions/>