

## ADAM17

### A Molecular Switch to Control TNFR2 During Atherogenesis In Vivo

Yiting Jia, Wei Kong

**A** disintegrin and metalloproteinase 17 (ADAM17) is a major sheddase of membrane-anchored cytokines, cell adhesion molecules, receptors, ligands, and enzymes, such as tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), transforming growth factor- $\alpha$ , TNF- $\alpha$  receptor 1 (TNFR1), TNF- $\alpha$  receptor 2 (TNFR2), and interleukin-6 receptor, L-selectin, Notch, and angiotensin-converting enzyme type 2, in which way ADAM17 could modulate a variety of signaling transduction and affect cell behavior.<sup>1-4</sup> The consequence of shedding events mainly depends on the type of substrate shed, the generation of soluble receptor agonists or antagonists, and the cell type and specific tissues affected during various disease states.

---

#### See accompanying article on page 247

---

ADAM17 was first identified to process membrane-bound TNF- $\alpha$  precursor to a soluble form. In addition, TNF receptors, TNFR1 and TNFR2, are both ADAM17 substrates. The potent cytokine TNF- $\alpha$  and TNF receptor signaling pathways control cell apoptosis, proliferation, adhesion, and inflammation and thus play an important role in pathogenesis and progression of several cardiovascular processes, including atherosclerosis.<sup>5,6</sup> TNFR1 is proved to be associated with aging-related coronary artery disease in humans, whereas TNFR2-deficient mice exhibit decreased plaque formation.<sup>7,8</sup> Interestingly, ADAM17 has been described to be upregulated in human and mouse atherosclerotic plaques,<sup>7,8</sup> and increased level of ADAM17 may contribute to atherosclerosis resistance,<sup>9,10</sup> which indicated a correlation between ADAM17, TNF signaling, and arteriosclerosis. Unfortunately, ADAM17 knockout mice are not viable because of its essential role in normal fetal development.<sup>11</sup> Mice with targeted deletion of exon 11 that encodes the catalytic active site of the metalloprotease domain (ADAM17 <sup>$\Delta$ Zn/ $\Delta$ Zn</sup>) also showed perinatal lethality within 2 weeks.<sup>12</sup> Then the conditional ADAM17 knockout animals have been generated to study the role of ADAM17 in physiological bone remodeling and neurological diseases.<sup>13</sup> However, the function of ADAM17 deficiency

in atherosclerosis is still in the mist. Recently, Chalaris et al generated partial deletion Adam 17<sup>ex/ex</sup> mice that showed only a weak ADAM17 expression in tissues, which may point the way to study the role of ADAM17 deficiency in atherosclerosis.<sup>14</sup>

In this issue of *Arteriosclerosis, Thrombosis, and Vascular Biology*, Nicolaou et al<sup>15</sup> crossed Adam 17<sup>ex/ex</sup> mice on Ldlr<sup>-/-</sup> background, then expanded on the previous studies of Holdt et al<sup>9</sup> and Canault et al<sup>10</sup> in Adam 17<sup>ex/ex</sup>Ldlr<sup>-/-</sup> mice. The authors revealed that atherosclerotic plaque size at the aortic root was markedly increased in Adam 17<sup>ex/ex</sup>Ldlr<sup>-/-</sup> mice compared with that of Adam 17<sup>wt/wt</sup>Ldlr<sup>-/-</sup> mice (1.5-folds), which indicated that ADAM17 deficiency would promote atherosclerosis. They found no alteration of total, LDL, and HDL cholesterol levels in ADAM17-deficient mice, which suggested that ADAM17 may affect atherosclerosis progression independent of plasma lipid level. Considering the known function of ADAM17 as a metalloproteinase, they systematically screened currently known ADAM17 substrates and consistently observed reduced release of TNFR1 and TNFR2, both in the plasma of Adam 17<sup>ex/ex</sup>Ldlr<sup>-/-</sup> mice and in the supernatant of Adam 17<sup>ex/ex</sup>Ldlr<sup>-/-</sup> bone marrow-derived macrophages. Subsequently, they detected increased level of TNF- $\alpha$  and TNFR2, but not TNFR1 on the cell membrane of Adam 17<sup>ex/ex</sup>Ldlr<sup>-/-</sup> macrophages compared with control macrophages, as well as elevated phosphorylation of bone marrow tyrosine kinase in chromosome X, a marker for constitutive activation of TNFR2 signaling. The proatherogenic effect of ADAM17 deficiency could be rescued by knockdown of TNFR2 in almost all of the vascular cells and macrophages detected. Finally, the authors detected less apoptotic cells and more proliferating macrophages in atherosclerotic lesions of Adam 17<sup>ex/ex</sup>Ldlr<sup>-/-</sup> mice, which confirmed the in vitro results. Taken together, these data provide convincing evidence that ADAM17 deficiency and TNFR2 signaling activation promote atherogenesis in mice (Figure).

This is the first definitive study of global ADAM17 knockout in atherosclerosis by loss of function mouse model in LDL<sup>-/-</sup> background. Although the authors examined ADAM17-TNFR2 signaling in almost all of the cell types involving plaque formation, we must realize that ADAM17 expression and activity varies in different cell types on proatherosclerotic stimuli. Previous studies by generating myeloid cell lineage-specific knockout ADAM17 mice revealed reduced proinflammatory cytokines secretion.<sup>14</sup> Endothelial-specific knockout ADAM17 mice exhibited reduced pathological neovascularization which may affect plaque vulnerability.<sup>16</sup> Considering ADAM17 has >70 different substrates, shedding by ADAM17 in different cell types needs to be individually

---

From the Department of Physiology and Pathophysiology, School of Basic Medical Sciences, Peking University, Beijing, China; and Key Laboratory of Molecular Cardiovascular Science, Ministry of Education, Beijing, China.

Correspondence to Wei Kong, PhD, Department of Physiology and Pathophysiology, Basic Medical College of Peking University, Beijing 100191, P.R. China. E-mail kongw@bjmu.edu.cn


(*Arterioscler Thromb Vasc Biol.* 2017;37:176-178.)

DOI: 10.1161/ATVBAHA.116.308840.)

© 2017 American Heart Association, Inc.

*Arterioscler Thromb Vasc Biol* is available at <http://atvb.ahajournals.org>

DOI: 10.1161/ATVBAHA.116.308840



**Figure. A**, Protective role of a disintegrin and metalloproteinase 17 (ADAM17) in atherosclerosis. **B**, ADAM17 deficiency reduces shedding of tumor necrosis factor receptor 2 (TNFR2) and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), thereby promotes membrane TNFR2 accumulation and persistent TNFR2 signaling activation, and enhances the downstream antiapoptosis and proproliferation effects in macrophages and vascular smooth muscle cells. BMX indicates bone marrow tyrosine kinase in chromosome X; sTNF- $\alpha$ , soluble TNF- $\alpha$ ; and sTNFR2, soluble TNFR2.

analyzed. Notably, recent development of novel proteomic methods may shed light on context-dependent bone fide substrates recognition.<sup>17</sup>

Overexpression of ADAM17 in atherosclerosis prone region raised the question how it is activated. Previous studies suggest that angiotensin II, reactive oxygen species, and polymethacrylate increase ADAM17 activity in multiple ways.<sup>18–20</sup> Inactive rhomboid types 1 and 2 have been reported to regulate forward trafficking of ADAM17 from endoplasmic reticulum to Golgi compartment where the prodomain of ADAM17 can be cleaved.<sup>21</sup> On the other hand, depletion of membrane cholesterol can induce ADAM17-dependent shedding.<sup>22</sup> More attention needs to be focused on the come and go of this ADAM17-TNFR2 axis. Additional studies are warranted to definitively demonstrate the role of ADAM17 in different cells and underlying regulatory mechanisms.

### Sources of Funding

This work was supported by funding from the National Natural Science Foundation of the P.R. China (NSFC; 91539203); International Cooperation and Exchanges NSFC (81220108004); the 111 Project of Chinese Ministry of Education (No. B07001); and the National Science Fund for Distinguished Young Scholars (81225002).

### Disclosures

None.

### References

- Black RA, Rauch CT, Kozlosky CJ, et al. A metalloproteinase disintegrin that releases tumour-necrosis factor- $\alpha$  from cells. *Nature*. 1997;385:729–733. doi: 10.1038/385729a0.
- Scheller J, Chalaris A, Garbers C, Rose-John S. ADAM17: a molecular switch to control inflammation and tissue regeneration. *Trends Immunol*. 2011;32:380–387. doi: 10.1016/j.it.2011.05.005.
- Wang R, Ye X, Bhattacharya R, Boulbes DR, Fan F, Xia L, Ellis LM. A disintegrin and metalloproteinase domain 17 regulates colorectal cancer

- stem cells and chemosensitivity via Notch1 signaling. *Stem Cells Transl Med*. 2016;5:331–338. doi: 10.5966/sctm.2015-0168.
- Tsukerman P, Eisenstein EM, Chavkin M, et al. Cytokine secretion and NK cell activity in human ADAM17 deficiency. *Oncotarget*. 2015;6:44151–44160. doi: 10.18632/oncotarget.6629.
- Lusis AJ. Atherosclerosis. *Nature*. 2000;407:233–241. doi: 10.1038/35025203.
- Pimentel-Muñoz FX, Seed B. Regulated commitment of TNF receptor signaling: a molecular switch for death or activation. *Immunity*. 1999;11:783–793.
- Chandrasekharan UM, Mavrakis L, Bonfield TL, Smith JD, DiCorleto PE. Decreased atherosclerosis in mice deficient in tumor necrosis factor- $\alpha$  receptor-II (p75). *Arterioscler Thromb Vasc Biol*. 2007;27:e16–e17. doi: 10.1161/01.ATV.0000255551.33365.22.
- Zhang L, Connelly JJ, Peppel K, Brian L, Shah SH, Nelson S, Crosslin DR, Wang T, Allen A, Kraus WE, Gregory SG, Hauser ER, Freedman NJ. Aging-related atherosclerosis is exacerbated by arterial expression of tumor necrosis factor receptor-1: evidence from mouse models and human association studies. *Hum Mol Genet*. 2010;19:2754–2766. doi: 10.1093/hmg/ddq172.
- Holdt LM, Thiery J, Breslow JL, Teupser D. Increased ADAM17 mRNA expression and activity is associated with atherosclerosis resistance in LDL-receptor deficient mice. *Arterioscler Thromb Vasc Biol*. 2008;28:1097–1103. doi: 10.1161/ATVBAHA.108.165654.
- Canault M, Peiretti F, Kopp F, Bonardo B, Bonzi MF, Coudeyre JC, Alessi MC, Juhan-Vague I, Nalbhone G. The TNF  $\alpha$  converting enzyme (TACE/ADAM17) is expressed in the atherosclerotic lesions of apolipoprotein E-deficient mice: possible contribution to elevated plasma levels of soluble TNF  $\alpha$  receptors. *Atherosclerosis*. 2006;187:82–91. doi: 10.1016/j.atherosclerosis.2005.08.031.
- Peschon JJ, Slack JL, Reddy P, et al. An essential role for ectodomain shedding in mammalian development. *Science*. 1998;282:1281–1284.
- Horiuchi K, Kimura T, Miyamoto T, Takaishi H, Okada Y, Toyama Y, Blobel CP. Cutting edge: TNF- $\alpha$ -converting enzyme (TACE/ADAM17) inactivation in mouse myeloid cells prevents lethality from endotoxin shock. *J Immunol*. 2007;179:2686–2689.
- Horiuchi K, Kimura T, Miyamoto T, Miyamoto K, Akiyama H, Takaishi H, Morioka H, Nakamura T, Okada Y, Blobel CP, Toyama Y. Conditional inactivation of TACE by a Sox9 promoter leads to osteoporosis and increased granulopoiesis via dysregulation of IL-17 and G-CSF. *J Immunol*. 2009;182:2093–2101. doi: 10.4049/jimmunol.0802491.
- Chalaris A, Adam N, Sina C, et al. Critical role of the disintegrin metalloprotease ADAM17 for intestinal inflammation and regeneration in mice. *J Exp Med*. 2010;207:1617–1624. doi: 10.1084/jem.20092366.

15. Nicolaou A, Zhao Z, Northoff BH, Sass K, Herbst A, Kohlmaier A, Chalaris A, Wolfrum C, Weber C, Steffens S, Rose-John S, Teupser D, Holdt LM. *Adam17* deficiency promotes atherosclerosis by enhanced TNFR2 signaling in mice. *Arterioscler Thromb Vasc Biol*. 2017;37:247–257. doi: 10.1161/ATVBAHA.116.308682.
16. Weskamp G, Mendelson K, Swendeman S, Le Gall S, Ma Y, Lyman S, Hinoki A, Eguchi S, Guaiquil V, Horiuchi K, Blobel CP. Pathological neovascularization is reduced by inactivation of ADAM17 in endothelial cells but not in pericytes. *Circ Res*. 2010;106:932–940. doi: 10.1161/CIRCRESAHA.109.207415.
17. Müller SA, Scilabra SD, Lichtenthaler SF. Proteomic substrate identification for membrane proteases in the brain. *Front Mol Neurosci*. 2016;9:96. doi: 10.3389/fnmol.2016.00096.
18. Lautrette A, Li S, Alili R, Sunnarborg SW, Burtin M, Lee DC, Friedlander G, Terzi F. Angiotensin II and EGF receptor cross-talk in chronic kidney diseases: a new therapeutic approach. *Nat Med*. 2005;11:867–874. doi: 10.1038/nm1275.
19. Myers TJ, Brennaman LH, Stevenson M, Higashiyama S, Russell WE, Lee DC, Sunnarborg SW. Mitochondrial reactive oxygen species mediate GPCR-induced TACE/ADAM17-dependent transforming growth factor- $\alpha$  shedding. *Mol Biol Cell*. 2009;20:5236–5249. doi: 10.1091/mbc.E08-12-1256.
20. Schwarz J, Broder C, Helmstetter A, Schmidt S, Yan I, Müller M, Schmidt-Arras D, Becker-Pauly C, Koch-Nolte F, Mittrücker HW, Rabe B, Rose-John S, Chalaris A. Short-term TNF $\alpha$  shedding is independent of cytoplasmic phosphorylation or furin cleavage of ADAM17. *Biochim Biophys Acta*. 2013;1833:3355–3367. doi: 10.1016/j.bbamcr.2013.10.005.
21. Li X, Maretzky T, Weskamp G, Monette S, Qing X, Issuree PD, Crawford HC, McIlwain DR, Mak TW, Salmon JE, Blobel CP. iRhoms 1 and 2 are essential upstream regulators of ADAM17-dependent EGFR signaling. *Proc Natl Acad Sci USA*. 2015;112:6080–6085. doi: 10.1073/pnas.1505649112.
22. Tellier E, Canault M, Rebsomen L, Bonardo B, Juhan-Vague I, Nalbone G, Peiretti F. The shedding activity of ADAM17 is sequestered in lipid rafts. *Exp Cell Res*. 2006;312:3969–3980. doi: 10.1016/j.yexcr.2006.08.027.

---

KEY WORDS: Editorials ■ apoptosis ■ atherosclerosis ■ bone marrow ■ cholesterol ■ ligands

# Arteriosclerosis, Thrombosis, and Vascular Biology



JOURNAL OF THE AMERICAN HEART ASSOCIATION

## ADAM17: A Molecular Switch to Control TNFR2 During Atherogenesis In Vivo Yiting Jia and Wei Kong

*Arterioscler Thromb Vasc Biol.* 2017;37:176-178

doi: 10.1161/ATVBAHA.116.308840

*Arteriosclerosis, Thrombosis, and Vascular Biology* is published by the American Heart Association, 7272  
Greenville Avenue, Dallas, TX 75231

Copyright © 2017 American Heart Association, Inc. All rights reserved.

Print ISSN: 1079-5642. Online ISSN: 1524-4636

The online version of this article, along with updated information and services, is located on the  
World Wide Web at:

<http://atvb.ahajournals.org/content/37/2/176>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Arteriosclerosis, Thrombosis, and Vascular Biology* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:

<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Arteriosclerosis, Thrombosis, and Vascular Biology* is online at:

<http://atvb.ahajournals.org/subscriptions/>