

## Oxidized Lipid Uptake by Scavenger Receptor CD36 (Cluster of Differentiation 36) Modulates Endothelial Surface Properties and May Contribute to Atherogenesis

Roy L. Silverstein

For many years, vascular biologists and pathologists have known that atherosclerosis is a spatially discontinuous disease; plaque tends to form in specific locations, especially branch points in large arteries, where blood flow is discontinuous and disordered. Furthermore, plaque formation progresses temporally from the proximal arterial tree to more distal vessels. It is also well known that vascular endothelial cells are sensitive to shear stress patterns and blood flow<sup>1</sup> and, in fact, express a robust mechanosensory system made up of intracellular signaling pathways that are triggered by integrins, selectins, and cilia.<sup>2</sup> Unidirectional flow in straight vessels, such as the distal aorta, is atheroprotective, whereas disordered flow (DF) at curvatures and branch points, such as the aortic arch, promotes inflammatory signaling, endothelial dysfunction, leukocyte recruitment, and plaque formation.<sup>3</sup> An article by Le Master et al<sup>4</sup> published in this issue of *Arteriosclerosis, Thrombosis, and Vascular Biology* presents elegant new data that helps integrate these observations into a pathogenic model that relates disordered blood flow to upregulated endothelial cell expression of the scavenger receptor CD36 (cluster of differentiation 36), increased uptake of oxidized lipids, and increased endothelial cell stiffness.

### See accompanying article on page 64

The team used atomic force microscopy, a powerful microscopy-based technique that uses a cantilevered probe to scan surfaces. With atomic force microscopy, they were able to measure endothelial monolayer stiffness in fresh aortae that were removed from mice and opened en face to expose the luminal surface. By measuring microindentation depth at multiple sites, a mean elastic modulus was calculated to estimate the degree of deformability or stiffness of the cell surface. Under basal conditions, they found that endothelial stiffness was  $\approx$ 2-fold greater in the atherosclerosis-prone aortic arch, compared with the distal aorta. Then, using a well-accepted model of moderate hyperlipidemia induced by feeding mice a high-fat diet for 1 month, they found a further 4-fold increased stiffness in the aortic arch from the hyperlipidemic mice. This was associated with a

significant increase in lipid and oxylipid accumulation in the aortic arch, suggesting possible mechanistic connections between DF, lipid accumulation, endothelial cell (EC) stiffness, and susceptibility to atherosclerosis.<sup>4</sup> To probe mechanisms, they also studied human aortic EC in culture using 2 well-characterized in vitro model systems of DF and showed that human aortic EC exposed to DF compared with laminar flow had increased uptake of oxidized lipids when incubated with oxLDL (oxidized low-density lipoprotein) and that this was associated with increased EC stiffness measured by atomic force microscopy.

Although oxLDL uptake and signaling through CD36 are well described in macrophages and clearly play critically important roles in foam cell formation, reactive oxygen species generation, inflammatory signaling, dysregulated cell migration, and plaque formation,<sup>5</sup> a role for this system in arterial EC is not well understood. CD36 is expressed prominently in microvascular EC where it mediates antiangiogenic, proapoptotic signaling,<sup>6</sup> but expression of CD36 in large-vessel EC has been controversial, and other scavenger receptors, such as LOX1 (lectin-like oxLDL receptor-1) and SRB1 (scavenger receptor B1),<sup>7</sup> have been proposed to be more important in endothelial atherogenesis. Le Master et al now show convincingly, using functional knockdown, mRNA expression assays, immunoblots, and immunofluorescence imaging, that human aortic EC in culture and mouse aortae in vivo express CD36. Although baseline expression is low ( $\approx$ 25% of that seen in capillary EC), it was increased by exposure to disturbed flow or oxLDL in vitro and by high-fat diet in vivo. Interestingly, CD36 expression was higher in the aortic arch compared with the distal aorta, consistent with a potential role in mediating the spatial differences in lipid uptake and endothelial stiffness seen in these 2 sites.

To probe these mechanistic connections, the authors used siRNA to knockdown CD36 expression in cultured human aortic EC and studied aortae isolated from *cd36*-null mice. They found that loss of CD36 abrogated the in vitro effects of DF plus oxLDL on both oxLDL uptake and increased stiffness. Importantly, the *cd36*-null mice on high-fat diet had less lipid deposition in their aortic arches and did not show the increase in endothelial stiffness seen in wild-type animals. Transplant of *cd36*-null bone marrows into irradiated wild-type mice did not rescue the phenotype, and transplant of wild-type bone marrows into *cd36*-null mice did not recapitulate the phenotype, strongly suggesting that the observed effects of high-fat diet on aorta lipid uptake and monolayer stiffness were mediated by CD36 in the vasculature, not in monocytes or macrophages. Although the evidence points to a role for EC CD36 in promoting endothelial

From the Department of Medicine, Medical College of Wisconsin, and Blood Research Institute, Blood Center of Wisconsin, Milwaukee.

Correspondence to Roy L. Silverstein, MD, Department of Medicine, Medical College of Wisconsin, 9200 W Wisconsin Ave, Milwaukee, WI 53226. E-mail rsilverstein@mcw.edu

(*Arterioscler Thromb Vasc Biol.* 2018;38:4-5.  
DOI: 10.1161/ATVBAHA.117.310351.)

© 2017 American Heart Association, Inc.

*Arterioscler Thromb Vasc Biol* is available at <http://atvb.ahajournals.org>  
DOI: 10.1161/ATVBAHA.117.310351

stiffness, they cannot rule out roles for smooth muscle cells<sup>8</sup> or other CD36-expressing vascular cells in the process.

Left unanswered by this sophisticated work is just what is the biological and pathological relevance of endothelial stiffness? Does it contribute to vascular stiffness, which is generally thought to be mediated by changes in the extracellular matrix, but which is known to increase with aging<sup>9</sup> and with atherosclerosis progression, and to associate with endothelial dysfunction? Does it influence atherogenic processes, such as leukocyte transmigration, endothelial barrier function, or proinflammatory signaling? Is it related to plasma membrane cholesterol or oxysterol content and could it therefore influence membrane microdomain formation, membrane protein localization, and signal transduction? These are interesting questions for further study, but enthusiasm for the work must be tempered by published results,<sup>10</sup> showing that transplant of *apoe;cd36* double null bone marrow into *apoe*-null mice led to >80% reduction in aortae atherosclerosis area compared with *apoe*-null mice transplanted with *apoe*-null;*cd36* wild-type marrows. These data suggest strongly that it is CD36 expression on hematopoietic cells, rather than vascular cells, that accounts for atheroprotection seen by deleting CD36 in mouse models and may imply that CD36-mediated changes in aorta endothelial stiffness and lipid uptake may not directly impact atherogenesis.

### Sources of Funding

This work was supported by National Institutes of Health P01HL087018.

### Disclosures

None.

### References

1. DePaola N, Gimbrone MA Jr, Davies PF, Dewey CF Jr. Vascular endothelium responds to fluid shear stress gradients. *Arterioscler Thromb.* 1992;12:1254–1257.
2. Dai G, Kaazempur-Mofrad MR, Natarajan S, Zhang Y, Vaughn S, Blackman BR, Kamm RD, García-Cardena G, Gimbrone MA Jr. Distinct endothelial phenotypes evoked by arterial waveforms derived from atherosclerosis-susceptible and -resistant regions of human vasculature. *Proc Natl Acad Sci USA.* 2004;101:14871–14876. doi: 10.1073/pnas.0406073101.
3. Hahn C, Schwartz MA. Mechanotransduction in vascular physiology and atherogenesis. *Nat Rev Mol Cell Biol.* 2009;10:53–62. doi: 10.1038/nrm2596.
4. Le Master E, Huang R-T, Zhang C, et al. Proatherogenic flow increases endothelial stiffness via enhanced CD36-mediated oxidized low-density lipoproteins. *Arterioscler Thromb Vasc Biol.* 2018;38:64–75. doi: 10.1161/ATVBAHA.117.309907.
5. Chen Y, Kennedy DJ, Ramakrishnan DP, Yang M, Huang W, Li Z, Xie Z, Silverstein RL. CD36 mediates macrophage pro-atherogenic responses to oxidized LDL by recruiting Na/K-ATPase/Lyn complex. *Sci Signal.* 2015;8:ra91.
6. Silverstein RL, Febbraio M. CD36, a scavenger receptor involved in immunity, metabolism, angiogenesis, and behavior. *Sci Signal.* 2009;2:re3. doi: 10.1126/scisignal.272re3.
7. Cominacini L, Pasini AF, Garbin U, Davoli A, Tosetti ML, Campagnola M, Rigoni A, Pastorino AM, Lo Cascio V, Sawamura T. Oxidized low density lipoprotein (ox-LDL) binding to ox-LDL receptor-1 in endothelial cells induces the activation of NF-kappaB through an increased production of intracellular reactive oxygen species. *J Biol Chem.* 2000;275:12633–12638.
8. Li W, Febbraio M, Reddy SP, Yu DY, Yamamoto M, Silverstein RL. CD36 participates in a signaling pathway that regulates ROS formation in murine VSMCs. *J Clin Invest.* 2010;120:3996–4006. doi: 10.1172/JCI42823.
9. Kohn JC, Lampi MC, Reinhart-King CA. Age-related vascular stiffening: causes and consequences. *Front Genet.* 2015;6:112. doi: 10.3389/fgene.2015.00112.
10. Febbraio M, Guy E, Silverstein RL. Stem cell transplantation reveals that absence of macrophage CD36 is protective against atherosclerosis. *Arterioscler Thromb Vasc Biol.* 2004;24:2333–2338. doi: 10.1161/01.ATV.0000148007.06370.68.

# Arteriosclerosis, Thrombosis, and Vascular Biology



JOURNAL OF THE AMERICAN HEART ASSOCIATION

## Oxidized Lipid Uptake by Scavenger Receptor CD36 (Cluster of Differentiation 36) Modulates Endothelial Surface Properties and May Contribute to Atherogenesis Roy L. Silverstein

*Arterioscler Thromb Vasc Biol.* 2018;38:4-5

doi: 10.1161/ATVBAHA.117.310351

*Arteriosclerosis, Thrombosis, and Vascular Biology* is published by the American Heart Association, 7272  
Greenville Avenue, Dallas, TX 75231

Copyright © 2017 American Heart Association, Inc. All rights reserved.

Print ISSN: 1079-5642. Online ISSN: 1524-4636

The online version of this article, along with updated information and services, is located on the  
World Wide Web at:

<http://atvb.ahajournals.org/content/38/1/4>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Arteriosclerosis, Thrombosis, and Vascular Biology* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Arteriosclerosis, Thrombosis, and Vascular Biology* is online at:  
<http://atvb.ahajournals.org/subscriptions/>