

## TRPing out Platelet Calcium

### TRPM7 (Transient Receptor Potential Melastatin-Like 7) Modulates Calcium Mobilization and Platelet Function via Phospholipase C Interactions

Anh T.P. Ngo, Owen J.T. McCarty, Joseph E. Aslan

Originally identified as coagulation factor IV, calcium ( $\text{Ca}^{2+}$ ) is now well established as a key cofactor in the formation of the tenase and prothrombinase complexes on the extracellular surfaces of activated platelets to ultimately mediate fibrin generation and hemostasis.<sup>1</sup> Similarly,  $\text{Ca}^{2+}$  has long been known to serve an important intracellular role in orchestrating the cell biological responses of platelets in hemostatic plug formation.<sup>2-5</sup> During the past several decades, as biochemical efforts have identified and refined roles for  $\text{Ca}^{2+}$  as an essential second messenger in virtually all cells,<sup>3</sup> complementary studies of platelets have similarly detailed how spatiotemporal changes in intracellular  $\text{Ca}^{2+}$  levels regulate platelet granule secretion, cytoskeletal dynamics, aggregation, and other cell biological outputs underlying platelet physiology. On a general mechanistic level, changes in intracellular  $\text{Ca}^{2+}$  concentrations that trigger the cellular responses driving platelet function are solicited downstream of a variety of receptors that differentially activate phospholipase C (PLC) family members, resulting in inositol-1,4,5-trisphosphate ( $\text{IP}_3$ ) production and  $\text{IP}_3$  ( $\text{IP}_3$  receptor)-mediated release of  $\text{Ca}^{2+}$  from intracellular stores (Figure).<sup>4,6-9</sup>

#### See accompanying article on page 344

Given the critical importance of intracellular  $\text{Ca}^{2+}$  concentration to platelet function, diverse endeavors have aimed to understand and target the myriad of molecular processes regulating and driven by  $\text{Ca}^{2+}$  dynamics. In addition to mobilization from internal stores after  $\text{IP}_3$  generation, extracellular  $\text{Ca}^{2+}$  also enters platelets via store- and receptor-operated calcium entry routes involving more recently described players such as *Orai1*, *STIM1* (stromal interaction molecule 1), and transient receptor potential (TRP) family channels.<sup>3</sup> In this issue of *Arteriosclerosis, Thrombosis, and Vascular Biology*, Gotru et al<sup>10</sup> now uncover a novel mechanism by which the TRP subfamily member TRPM7 (transient receptor potential melastatin-like 7) modulates PLC phosphorylation and intracellular calcium mobilization to effect store-operated calcium

entry and platelet function. By taking advantage of a transgenic mouse model with a loss-of-function mutation in the cytosolic TRPM7 C-terminal serine/threonine kinase domain (*Trpm7<sup>RR</sup>*),<sup>11</sup> this study specifies a role for the TRPM7 kinase domain—rather than its constitutive  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$  channel activity—in modulating the phosphorylation and activation of PLC family members and consequently intracellular calcium responses and platelet functional processes underlying both hemostasis and thrombosis (Figure).

Earlier work from Stritt et al<sup>12</sup> suggested a general role for TRPM7 in regulating platelet function through pathways controlling megakaryocyte and platelet cytoskeletal dynamics. In the current study, Gotru et al<sup>10</sup> now more specifically show altered signaling responses in *Trpm7<sup>RR</sup>* platelets, including a delay in phosphorylation of Syk (spleen tyrosine kinase), LAT (linker for activation of T-cells),  $\text{PLC}\gamma 2$ , and  $\text{PKC}\epsilon$  (protein kinase C) in response to the platelet GPVI (glycoprotein VI) receptor agonist collagen-related peptide. Similar effects are also found to be associated with rhodocytin→CLEC-2 (C-type lectin-like receptor 2)→ $\text{PLC}\gamma 2$  as well as thrombin→PAR (protease-activated receptor)→ $\text{PLC}\beta 3$  signaling axes. To investigate the physiological consequences associated with these alterations in signaling kinetics in platelets, Gotru et al also analyze many platelet phenotypes and functional responses in *Trpm7<sup>RR</sup>* mice, revealing defects in granule secretion, integrin activation, and platelet aggregation *in vitro*. In association with these altered platelet responses, *Trpm7<sup>RR</sup>* mice display prolonged tail bleeding times and significantly less vessel occlusion in response to arteriole injury compared with wild-type counterparts, supporting roles for the TRPM7 kinase domain in modulating platelet function *in vivo*. Perhaps most remarkably, antithrombotic effects of mutating the TRPM7 kinase domain seem to be mediated through marrow-derived cells, especially platelets, because wild-type chimeric mice transplanted with bone marrow from *Trpm7<sup>RR</sup>* mice, or thrombocytopenic wild-type mice transfused with platelets from *Trpm7<sup>RR</sup>* animals, were similarly protected from infarct progression and showed overall improved neurological and motor function outcomes relative to matched controls after transient middle cerebral artery occlusion to model ischemic stroke.

Although Gotru et al have astutely elucidated platelet-associated roles for the TRPM7 kinase domain in complete and proper platelet function, many intriguing questions remain on how TRPM7 impacts platelet physiology in hemostasis and thrombosis. For instance, it is not clear how mutation of this serine/threonine kinase domain translates into a delay in the tyrosine phosphorylation of receptor-proximal components of platelet-signaling systems, especially as physiological substrates of TRPM7 are limited and have not yet


From the Department of Biomedical Engineering (A.T.P.N., O.J.T.M.), Department of Biochemistry and Molecular Biology (J.E.A.), and Knight Cardiovascular Institute (J.E.A.), School of Medicine, Oregon Health & Science University, Portland.

Correspondence to Joseph E. Aslan, PhD, 3181 SW Sam Jackson Park Rd, Mail Code HRC5N, Oregon Health & Science University, Portland, OR 97239. E-mail aslanj@ohsu.edu

(*Arterioscler Thromb Vasc Biol.* 2018;38:285-286.  
DOI: 10.1161/ATVBAHA.117.310493.)

© 2018 American Heart Association, Inc.

*Arterioscler Thromb Vasc Biol* is available at <http://atvb.ahajournals.org>  
DOI: 10.1161/ATVBAHA.117.310493



**Figure.** TRPM7 (transient receptor potential melastatin-like 7) kinase domain interacts with phospholipase C (PLC) family members to regulate intracellular  $\text{Ca}^{2+}$  responses and platelet function (model). Platelet activation downstream of GPVI (glycoprotein VI), CLEC-2 (C-type lectin-like receptor 2), PARs (protease-activated receptors), and other receptors results in intracellular signaling events that upregulate PLC (PLC $\gamma$ 2 and PLC $\beta$ 3) phosphorylation and activity to drive the metabolism of phosphatidylinositol 4,5-bisphosphate ( $\text{PIP}_2$ ) on the cytosolic face of the platelet plasma membrane to produce inositol-1,4,5-trisphosphate ( $\text{IP}_3$ ) and diacylglycerol (DAG). Rather than serving as a constitutively active  $\text{Mg}^{2+}/\text{Ca}^{2+}$  channel or regulating DAG-mediated receptor-operated calcium entry (ROCE) through TRP channels (TRPC) and other associated processes, this study supports a model whereby the TRPM7 kinase domain interacts with PLC family members to modulate PLC phosphorylation and activation,  $\text{IP}_3$  production, intracellular  $\text{Ca}^{2+}$  mobilization, and STIM1-Orai1-mediated store-operated calcium entry (SOCE), ultimately modulating intracellular  $\text{Ca}^{2+}$  concentrations and associated  $\text{Ca}^{2+}$  responses underlying platelet function.<sup>10</sup> Illustration provided by Inky Mouse Studios ©2018—all rights reserved.

been demonstrated in platelets. Furthermore, these findings remain to be placed into the context of the former studies by Stritt et al,<sup>12</sup> which made use of megakaryocyte/platelet lineage-specific deletions of TRPM7 that resulted in altered cytoskeletal phenotypes. Along these lines, as platelets from *Trpm7<sup>R/R</sup>* mice exhibit delayed PKC phosphorylation kinetics, many other signaling processes around the platelet cytoskeleton may be disrupted, especially given the spatial and temporal link between intracellular calcium mobilization and PKC activation in regulating Rho GTPases and platelet function.<sup>13,14</sup> Finally, from a translational perspective, the rather unique TRPM7 kinase domain may represent a specialized therapeutic or biomarker target; however, validation of the relevance of TRPM7 within human platelet biology remains to be shown. Regardless, whether TRPM7 serves as a kinase, a molecular scaffold, an effector, or some combination thereof, Gotru et al provide a rich contribution to the ever-evolving and increasingly important understanding of mechanisms regulating intracellular  $\text{Ca}^{2+}$  dynamics in health and disease in platelet physiology and beyond.

### Sources of Funding

We are supported by grants from the National Institutes of Health (R01HL101972 and R01GM116184 to O.J.T. McCarty) and the American Heart Association (17SDG33350075 to J.E. Aslan and 13EIA12630000 to O.J.T. McCarty).

### References

1. Wolberg AS. Thrombin generation and fibrin clot structure. *Blood Rev.* 2007;21:131–142. doi: 10.1016/j.blre.2006.11.001.
2. Wright JH, Minot GR. The viscous metamorphosis of the blood platelets. *J Exp Med.* 1917;26:395–409.
3. Varga-Szabo D, Braun A, Nieswandt B. Calcium signaling in platelets. *J Thromb Haemost.* 2009;7:1057–1066. doi: 10.1111/j.1538-7836.2009.03455.x.
4. Li Z, Delaney MK, O'Brien KA, Du X. Signaling during platelet adhesion and activation. *Arterioscler Thromb Vasc Biol.* 2010;30:2341–2349. doi: 10.1161/ATVBAHA.110.207522.
5. Aslan JE. Platelet shape change. In: Greslele P, Kleiman NS, Lopez JA, Page CP, eds. *Platelets in Thrombotic and Non-Thrombotic Disorders: Pathophysiology, Pharmacology and Therapeutics: An Update.* Cham: Springer International Publishing; 2017:321–336.
6. Feinstein MB, Fraser C. Human platelet secretion and aggregation induced by calcium ionophores. Inhibition by PGE1 and dibutyryl cyclic AMP. *J Gen Physiol.* 1975;66:561–581.
7. Wallach DF, Surgenor DM, Steele BB. Calcium-lipid complexes in human platelets. *Blood.* 1958;13:589–598.
8. Hathaway DR, Adelstein RS. Human platelet myosin light chain kinase requires the calcium-binding protein calmodulin for activity. *Proc Natl Acad Sci USA.* 1979;76:1653–1657.
9. McCarty OJ, Zhao Y, Andrew N, Machesky LM, Staunton D, Frampton J, Watson SP. Evaluation of the role of platelet integrins in fibronectin-dependent spreading and adhesion. *J Thromb Haemost.* 2004;2:1823–1833. doi: 10.1111/j.1538-7836.2004.00925.x.
10. Gotru SK, Chen W, Kraft P, et al. TRPM7 kinase controls calcium responses in arterial thrombosis and stroke in mice. *Arterioscler Thromb Vasc Biol.* 2018;38:344–352. doi: 10.1161/ATVBAHA.117.310391.
11. Matsushita M, Kozak JA, Shimizu Y, McLachlin DT, Yamaguchi H, Wei FY, Tomizawa K, Matsui H, Chait BT, Cahalan MD, Nairn AC. Channel function is dissociated from the intrinsic kinase activity and autophosphorylation of TRPM7/ChaK1. *J Biol Chem.* 2005;280:20793–20803. doi: 10.1074/jbc.M413671200.
12. Stritt S, Nurden P, Favier R, et al. Defects in TRPM7 channel function deregulate thrombopoiesis through altered cellular  $\text{Mg}^{2+}$  homeostasis and cytoskeletal architecture. *Nat Commun.* 2016;7:11097. doi: 10.1038/ncomms11097.
13. Aslan JE, McCarty OJ. Rho GTPases in platelet function. *J Thromb Haemost.* 2013;11:35–46. doi: 10.1111/jth.12051.
14. Ngo AT, Thierheimer ML, Babur Ö, Rocheleau AD, Huang T, Pang J, Rigg RA, Mitrugno A, Theodorescu D, Burchard J, Nan X, Demir E, McCarty OJ, Aslan JE. Assessment of roles for the Rho-specific guanine nucleotide dissociation inhibitor Ly-GDI in platelet function: a spatial systems approach. *Am J Physiol Cell Physiol.* 2017;312:C527–C536. doi: 10.1152/ajpcell.00274.2016.

# Arteriosclerosis, Thrombosis, and Vascular Biology



JOURNAL OF THE AMERICAN HEART ASSOCIATION

## TRPing out Platelet Calcium: TRPM7 (Transient Receptor Potential Melastatin-Like 7) Modulates Calcium Mobilization and Platelet Function via Phospholipase C Interactions

Anh T.P. Ngo, Owen J.T. McCarty and Joseph E. Aslan

*Arterioscler Thromb Vasc Biol.* 2018;38:285-286

doi: 10.1161/ATVBAHA.117.310493

*Arteriosclerosis, Thrombosis, and Vascular Biology* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2018 American Heart Association, Inc. All rights reserved.

Print ISSN: 1079-5642. Online ISSN: 1524-4636

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://atvb.ahajournals.org/content/38/2/285>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Arteriosclerosis, Thrombosis, and Vascular Biology* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:

<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Arteriosclerosis, Thrombosis, and Vascular Biology* is online at:

<http://atvb.ahajournals.org/subscriptions/>